M. Strosznajder Songin. Glycogen synthase kinase 3beta and its phosphorylated form (Y216) in the paraquat-induced m S. K. McCormack Mak. Lysosomal degradation of alpha-synuclein in vivo. J Biol Chem. 2010. 285:13621-9 M. J. Gil Kang. Selective vulnerability of the striatal subregions of C57BL/6 mice to paraquat. Toxicol Lett. 2010. 195:127 W. S. Abel Choi. JNK3 mediates paraquat- and rotenone-induced dopaminergic neuron death. J Neuropathol Exp Neuro H. O. Chang Lawal. The Drosophila vesicular monoamine transporter reduces pesticide-induced loss of dopaminergic ne A. Ahmad Kumar. Effect of zinc and paraguat co-exposure on neurodegeneration: Modulation of oxidative stress and ex L. Lu Yin. Genetic-based, differential susceptibility to paraquat neurotoxicity in mice. Neurotoxicol Teratol. 2011. 33:41 E. N. Peters Mangano. Granulocyte macrophage-colony stimulating factor protects against substantia nigra dopaminerg R. Sanchez Bajo-Graneras. Apolipoprotein D alters the early transcriptional response to oxidative stress in the adult cere E. N. Litteljohn Mangano. Interferon-gamma plays a role in paraquat-induced neurodegeneration involving oxidative an C. B. Lazo Hutson. Traumatic brain injury in adult rats causes progressive nigrostriatal dopaminergic cell loss and enhand R. Ganfornina Bajo-Graneras. Apolipoprotein D mediates autocrine protection of astrocytes and controls their reactivity G. Candiani Zega. Neurotoxic effect of the herbicide paraquat on ascidian larvae. Environ Toxicol Pharmacol. 2010. 29:2 A. Karadayian Czerniczyniec. Paraquat induces behavioral changes and cortical and striatal mitochondrial dysfunction. F D. Nelson Litteljohn. The effects of paraquat on regional brain neurotransmitter activity, hippocampal BDNF and behavi H. Huang Zhou. Early exposure to paraquat sensitizes dopaminergic neurons to subsequent silencing of PINK1 gene exp S. Chakrabarti Mitra. Differential regional expression patterns of alpha-synuclein, TNF-alpha, and IL-1beta; and variable P. M. Cui Rappold. Paraguat neurotoxicity is mediated by the dopamine transporter and organic cation transporter-3. Pi A. Singh Kumar. Involvement of NADPH oxidase and glutathione in zinc-induced dopaminergic neurodegeneration in ra Y. Lu Jiao. Genetic dissection of strain dependent paraquat-induced neurodegeneration in the substantia nigra pars com G. Dixit Srivastava. Resveratrol potentiates cytochrome P450 2 d22-mediated neuroprotection in maneb- and paraquat-R. Rong Pattarini. Long-lasting transcriptional refractoriness triggered by a single exposure to 1-methyl-4-phenyl-1,2,3,6 J. Mangano Bobyn. Viral-toxin interactions and Parkinson's disease: poly I:C priming enhanced the neurodegenerative e H. Wu Li. Neuroprotective effects of tert-butylhydroquinone on paraquat-induced dopaminergic cell degeneration in C5 A. C. Guhathakurta Cristovao. NADPH oxidase 1 mediates alpha-synucleinopathy in Parkinson's disease. J Neurosci. 201 M. A. Shehab Fahim. Daily subacute paraquat exposure decreases muscle function and substantia nigra dopamine level M. B. Nobuta Watson. PACAP deficiency sensitizes nigrostriatal dopaminergic neurons to paraquat-induced damage and C. B. Sturgess Breckenridge. Pharmacokinetic, neurochemical, stereological and neuropathological studies on the poten S. K. Prakash Yaday. Mucuna pruriens seed extract reduces oxidative stress in nigrostriatal tissue and improves neurobe A. Lores-Arnaiz Czerniczyniec. Mitochondrial susceptibility in a model of paraquat neurotoxicity. Free Radic Res. 2013. L. P. Kavanagh Liang. Glutathione deficiency in Gclm null mice results in complex I inhibition and dopamine depletion fo F. L. Carvalho Campos. Rodent models of Parkinson's disease: beyond the motor symptomatology. Front Behav Neurosc D. J. Travis Minnema. Dietary administration of paraquat for 13 weeks does not result in a loss of dopaminergic neurons S. Tadros Nuber. Environmental neurotoxic challenge of conditional alpha-synuclein transgenic mice predicts a dopamir J. W. Cognato Bortolotto. Long-term exposure to paraquat alters behavioral parameters and dopamine levels in adult ze A. K. Pragya Shukla. Mutation in Drosophila methuselah resists paraquat induced Parkinson-like phenotypes. Neurobiol J. A. Hessner Navarro. Analysis of dopaminergic neuronal dysfunction in genetic and toxin-induced models of Parkinson R. Krishna Hosamani. Standardized Bacopa monnieri extract ameliorates acute paraquat-induced oxidative stress, and r L. Achumi Phom. Curcumin's neuroprotective efficacy in Drosophila model of idiopathic Parkinson's disease is phase spe C. P. Leung Gonzalez-Hunt. Exposure to mitochondrial genotoxins and dopaminergic neurodegeneration in Caenorhabd L Na Chen. MERPS inflammasome activation by mitochondrial reactive oxygen species plays a key role in long-term cos C. Walker Coughlan. Comparative Proteomic Analysis of Carbonylated Proteins from the Striatum and Cortex of Pesticid C. Litteljohn Rudyk. Paraquat and psychological stressor interactions as pertains to Parkinsonian co-morbidity. Neurobid C. Niu Su. Low doses of single or combined agrichemicals induces alpha-synuclein aggregation in nigrostriatal system of N. Froio Degori. BEHAVIORAL AND ELECTROCORTICAL CHANGES INDUCED BY PARAQUAT AFTER INJECTION IN SPECIFIC Jayshree P Nellore. Paraquat exposure induces behavioral deficits in larval zebrafish during the window of dopamine ne Q. Chen, Y. Niu, R. Zhang, H. Guo, Y. Gao, Y. Li, R. Liu. The toxic influence of paraguat on hippocampus of mice: involvem

Parkinson's disease is a slowly progressing disease, due to a lesion of dopaminergic neurons in the substantia nigra and a Pathologic accumulation of alpha-synuclein is a feature of human parkinsonism and other neurodegenerative diseases. T Paraguat (PQ) is a strong redox agent that contributes to the formation of reactive oxygen species (ROS) and induces toxi Mechanistic studies underlying dopaminergic neuron death may identify new drug targets for the treatment of Parkinsor Dopamine is cytotoxic and may play a role in the development of Parkinson's disease. However, its interaction with envir Oxidative stress is implicated in Parkinson's disease (PD). Metallothioneins (MT), cytochrome P450 IIE1 (CYP2E1) and glu Paraquat (PQ) is an herbicide used extensively in agriculture. This agent is also suspected to be a risk factor for Parkinson Parkinson's disease (PD) has been linked to exposure to a variety of chemical (e.g., pesticides) and inflammatory agents, The lipocalin Apolipoprotein D (ApoD), known to protect the nervous system against oxidative stress (OS) in model organ Exposure to environmental contaminants, particularly pesticides, may be an important etiological factor in Parkinson's di Parkinson's disease (PD) is a neurodegenerative disorder characterized by the loss of nigrostriatal dopaminergic neurons The study of glial derived factors induced by injury and degeneration is important to understand the nervous system resp Paraquat is an herbicide widely used in agriculture, that proved to have toxic effect on many animal models. Moreover, it Paraguat is a highly toxic quaternary nitrogen herbicide capable of increasing superoxide anion production. The aim of the Accumulating evidence implicates pesticides such as paraquat in the development of Parkinson's disease (PD). Indeed, pa Environmental exposure, genetic modification, and aging are considered risky for Parkinson's disease (PD). How these ris BACKGROUND: Paraquat (1, 1-dimethyl-4, 4-bipyridium dichloride; PQ) causes neurotoxicity, especially dopaminergic ne The herbicide paraquat (PQ) has increasingly been reported in epidemiological studies to enhance the risk of developing An association between excessive zinc (Zn) accumulation in brain and incidences of Parkinson's disease (PD) has been sh The etiology of the vast majority of Parkinson's disease (PD) cases is unknown. It is generally accepted that there is an int A strong association between polymorphisms of the cytochrome P450 (CYP/Cyp) 2D6 gene and risk to Parkinson's diseasc Parkinson's disease (PD) is a progressive neurodegenerative disorder whose etiology is thought to have environmental (t BACKGROUND: Parkinson's disease (PD) has been linked with exposure to a variety of environmental and immunological The present study was aimed at determining the role of paraquat (PQ) in the activation of the NF-E2-related factor 2 (Nrf Accumulation of misfolded alpha-synuclein is the pathological hallmark of Parkinson's disease (PD). Nevertheless, little is The use of the herbicide paraquat (1,1'-dimethyl-4,4'-bipyridylium dichloride; PQ) which is widely used in agriculture is k Exposure to the pesticide paraquat (PQ) increases the risk of Parkinson's disease (PD), and its effect may be modulated b The pharmacokinetics and neurotoxicity of paraquat dichloride (PQ) were assessed following once weekly administration Parkinson's disease (PD) is a neurodegenerative disease which causes rigidity, resting tremor and postural instability. Tre Paraquat is a highly toxic herbicide capable of generating oxidative stress and producing brain damage after chronic expo Depletion of glutathione has been shown to occur in autopsied brains of patients with Parkinson's disease (PD) and in ani Parkinson's disease (PD) is classically characterized by motor symptoms; however, non-motor symptoms (NMS) are incre Several investigations have reported that mice administered paraquat dichloride (PQ.Cl2) by intraperitoneal injection exl The olfactory bulb (OB) is one of the first brain regions in Parkinson's disease (PD) to contain alpha-synuclein (alpha-syn) Chronic exposure to paraquat (Pq), a toxic herbicide, can result in Parkinsonian symptoms. This study evaluated the effec Parkinson's disease (PD) is a prevalent and devastating neurodegenerative disorder having limited cure options and stro Drosophila melanogaster has contributed significantly to the understanding of disease mechanisms in Parkinson's disease Objectives Bacopa monnieri (BM), an ayurvedic medicinal plant, has attracted considerable interest owing to its diverse r Selective degeneration of dopaminergic neurons in the substantia nigra underlies the basic motor impairments of Parkin Neurodegeneration has been correlated with mitochondrial DNA (mtDNA) damage and exposure to environmental toxin

Epidemiological studies indicate exposures to the herbicide paraquat (PQ) and fungicide maneb (MB) are associated with A number of epidemiological and experimental studies have implicated the non-selective herbicide, paraquat, in the development (SNCA) genes and environmental factors are important risk factors of Parkinson's disease (PD). The agric AREAS OF THE BRAIN OF THE RAT. Neuropharmacology. 1988. 27:201-207

Exposure to environmental risk factors such as herbicides in early life has been proposed to play important roles in the d Environmental paraquat (PQ) exposure has been suggested to be a potential risk factor for neurodegenerative disorders

AniPrim	NOADME		Rats
his accumulatio			
			Mice
AniPrim	NOADME		Mice
AniPrim	NOADME		Mice
AniPrim	NOADME		Drosophila
AniPrim	NOADME	1	Rat
AniPrim	YESADME		Mice
AniPrim	NOADME		Mice
isms, is up-regu	NOADME		
AniPrim	NOADME		Mice
AniPrim	NOADME		Rats
AniPrim	NOADME		Mice
t is considered a	NOADME		P. mammillata, C. intestinalis
AniPrim	NOADME		Rats
AniPrim	NOADME		Mice
AniPrim	NOADME		Rats
AniPrim	NOADME		Mice
e (PD) is well es			Mice
oxin) and genet			Mice
AniPrim	NOADME		Mice
AniPrim	NOADME		Mice
AniPrim	NOADME	***************************************	Rats
AniPrim	NOADME		Rats
AniPrim	NOADME		Mice
AniPrim	YESADME		Mice
AniPrim	NOADME		Mice
sure. The aim o			Rats
mal models of f			Mice
AniPrim	NOADME		Rats
AniPrim		***************************************	Mice
	YESADME		
AniPrim	NOADME		Mice
AniPrim	NOADME		Zebrafish
AniPrim	YESADME	***************************************	Drosophila
e (PD) as it is on			Drosophila
neuropharmaco			Mice
AniPrim	NOADME		Drosophila
s, but causation			C. elegans
AmiPilm	NOACRAE		
AniPrim	NOADME		Mice
elopment of spo			Mice
AniPrim	NOADME		Mice
AniPrim	NOADME		Rats
AniPrim	NOADME		Zebrafish
AniPrim	NOADME		Mice

subchronic
subacute
subscuto
subacute
subacute
subacute
developmental/acute
subacute
subacute
acute, subacute
subacute
subacute
subacute
subacute
subacute
subacute
subacute
subacute
subacute
subacute
acute
acute, subacute
subacute
subacute
subacute
subacute
subchronic
subacute
subacute
subscute
subacute
acute
subacute
acute/subacute
subacute
subacute
subacute
acute
acute/subacute
subacute

37 weeks, once a week
3 days, one dosage
3 weeks, twice per week
6 weeks, twice per week
up to 60 days
2-8 weeks, twice per week
3 weeks, once per week
3 weeks, 3 times per week
3 weeks, 3 times per week
2 weeks, once a week
5 weeks, 2 times per week in first 2 weeks and once per week for next three weeks
about 18 hours
4 weeks, once per week
3 weeks, 3 times per week
once (14-day olds)3 weeks, 2 times per week (12-week olds)
4 weeks, twice a week
20 days, every two days
8 weeks, twice per week
3 weeks, twice per week
9 weeks, twice per week
4 days, once on day 0 and once on day 4
3 weeks, 3 times per week
10 doses, 2-day intervals
4 injections, separated by one day
3 weeks, daily
single dose
once, twice or thrice, each dose separated by a week
3, 6, and 9 weeks, twice per week
4 weeks, once per week
3 weeks, twice per week
28 days, once per day
13 weeks
6 weeks, twice per week
6 injections, once every 3 days
12 and 24 hours
up to 8 days
single dose
up to 13 days
96 hours
6 weeks, twice per week
6.5 weeks, twice per week
4 weeks, twice per week
single dose
up to 96 hours
28 days, once per day